

1 Expectation

Assume Ω is a sample space

$$\Omega = \{x_1, x_2, x_3, \dots, x_n\}$$

Assume p is a probability distribution over Ω . Assume f is a function from Ω to numbers:

$$f \in \mathbb{R}^\Omega$$

We define the **expected value of f** :

$$E(f) = p(x_1)f(x_1) + p(x_2)f(x_2) + p(x_3)f(x_3) + \dots + p(x_n)f(x_n)$$

$$E(f) = \sum p(x_i)f(x_i)$$

2 Entropy

We assume a random variable X defined on an alphabet of symbols χ with pmf p . So the kinds of events we are now interested in are symbol occurrences. And we assume that the information measure of each symbol x , $x \in \chi$ is:

$$I(x) = -\log p(x)$$

We have:

$$E(I) = \sum p(x)(-\log p(x)) = -\sum p(x) \log p(x)$$

We call $E(I)$ the **entropy of random variable X** . It is usually written H :

$$H(X) = -\sum_{x \in \chi} p(x) \log p(x)$$

3 Entropy of Joint and Conditional Distributions

The entropy of a joint distribution is the same as the entropy of a distribution of a single random variable, except that the elements of the sample space are pairs:

$$H(X, Y) = -\sum_{x \in \chi} p(x, y) \log p(x, y)$$

Conditional entropy is somewhat different:

$$H(Y|X) = -\sum_{x \in \chi} p(x) H(p(Y|X = x))$$

Recall that $p(Y|X)$ is not a real pmf. What we do is sum up the entropies of **each** of the conditional pmfs of the form $p(Y|X=x)$, weighting each by the probability of x

$$H(Y|X) = - \sum_{x \in \mathcal{X}} p(x) \left[- \sum_{y \in \mathcal{Y}} p(y|x) \log p(y|x) \right]$$

Multiplying $p(x)$ into the inner sum:

$$H(Y|X) = - \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x) p(y|x) \log p(y|x)$$

Using the Chain rule:

$$H(Y|X) = - \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log p(y|x)$$

It turns out this is a rather natural definition of $H(X|Y)$ because it leads to a chain rule for entropy (proof, p. 64 of text):

$$H(X, Y) = H(Y|X) + H(X)$$