1 Expectation

Assume Ω is a sample space

$$\Omega = \{x_1, x_2, x_3, \ldots x_n\}$$

Assume p is a probability distribution over Ω. Assume f is a function from Ω to numbers:

$$f \in \aleph^{\Omega}$$

We define the **expected value of f**:

$$\mathrm{E}(f) = p(x_1)f(x_1) + p(x_2)f(x_2) + p(x_3)f(x_3) + \ldots + p(x_n)f(x_n)$$

$$\mathrm{E}(f) = \sum p(x_i)f(x_i)$$

2 Entropy

We assume a random variable X defined on an alphabet of symbols χ with pmf p. So the kinds of events we are now interested in are symbol occurrences. And we assume that the information measure of each symbol x, $x \in \chi$ is:

$$\mathrm{I}(x) = -\log p(x)$$

We have:

$$E(I) = \sum p(x)(-\log p(x)) = -\sum p(x)\log p(x)$$

We call E(I) the **entropy of random variable X**. It usually written H:

$$\mathrm{H}(X) = -\sum_{x \in \chi} p(x)\log p(x)$$

3 Entropy of Joint and Conditional Distributions

The entropy of a joint distribution is the same as the entropy of a distribution of a single random variable, except that the elements of the sample space are pairs:

$$\mathrm{H}(X, Y) = -\sum_{x \in \chi} p(x, y)\log p(x, y)$$

Conditional entropy is somewhat different:

$$\mathrm{H}(Y|X) = -\sum_{x \in \chi} p(x)H(p(Y|X = x))$$

Recall that p(Y |X) is not a real pmf. What we do is sum up the entropies of **each** of the conditional pmfs of the form p(Y | X=x), weighting each by the probability of x

$$\mathrm{H}(Y|X) = -\sum_{x\in\chi} p(x)[-\sum_{y\in\chi} p(y|x)\log p(y|x)]$$

Multiplying p(x) into the inner sum:

$$\mathrm{H}(Y|X) = -\sum_{x\in\chi}\sum_{y\in\chi} p(x)p(y|x)\log p(y|x)$$

Using the Chain rule:

$$\mathrm{H}(Y|X) = -\sum_{x\in\chi}\sum_{y\in\chi} p(x,y)\log p(y|x)$$

It turns out this is a rather natural definition of H(X|Y) because it leads to a chain rule for entropy (proof, p. 64 of text):

$$\mathrm{H}(X,Y) = \mathrm{H}(Y|X) + \mathrm{H}(X)$$